Mon. Not. R. Astron. Soc. **000**, 1–6 (0000) Printed 23 June 2008 (MN LaTeX style file v2.2)

arXiv:0803.3944v3 [astro-ph] 23 Jun 2008

# Towards Understanding the Mass-Metallicity relation of Quasar Absorbers: Evidence for bimodality and consequences.

Nicolas Bouché
[1]*Max Planck Institut für extraterrestrische Physik, Giessenbachstrasse, Garching D-85748, Germany; nbouche@mpe.mpg.de*

Accepted 2008 June 7. Received 2008 May 29; in original form 2008 April 7

**ABSTRACT**
One way to characterize and understand H I-selected galaxies is to study their metallicity properties. In particular, we show that the metallicity of absorbers is a bivariate function of the H I column density ($N_{\rm H\,I}$) and the Mg II equivalent width ($W_{\rm r}^{\lambda2796}$). Thus, a selection upon $W_{\rm r}^{\lambda2796}$ is not equivalent to a H I selection for intervening absorbers. A direct consequence for damped absorbers with $\log N_{\rm H\,I} > 20.3$ that falls from the bivariate metallicity distribution is that any correlation between the metallicity [X/H] and velocity width (using $W_{\rm r}^{\lambda2796}$ as a proxy) cannot be interpreted as a signature of the mass-metallicity relation akin to normal field galaxies. In other words, DLA samples are intrinsically heterogeneous and the [X/H]–$W_{\rm r}^{\lambda2796}$ or [X/H]–$\Delta v$ correlation reported in the literature arises from the H I cut. On the other hand, a sample of Mg II-selected absorbers, which are statistically dominated by lowest $N_{\rm H\,I}$ systems (sub-DLAs) at each $W_{\rm r}^{\lambda2796}$, are found to have a more uniform metallicity distribution. We postulate that the bivariate distribution [[X/H]($N_{\rm H\,I}$,$W_{\rm r}^{\lambda2796}$)] can be explained by two different physical origins of absorbers, namely sight-lines through the ISM of small galaxies and sight-lines through out-flowing material. Several published results follow from the bivariate [X/H] distribution such as (a) the properties of the two classes of DLAs, reported by Wolfe et al., and (b) the constant dust-to-gas ratio for Mg II-absorbers.

**Key words:** cosmology: observations — galaxies: evolution — galaxies: halos — quasars: absorption lines

## 1 INTRODUCTION

Among quasar (QSO) absorption line systems, damped Ly$\alpha$ absorbers (DLAs) with column densities $N_{\rm H\,I} \geqslant 2\times10^{20}$ cm$^{-2}$ are the most puzzling. Their high column density range indicates that they ought to trace cold neutral gas, hence disks (e.g. Wolfe et al. 1986; Wolfe & Prochaska 1998). This is further supported as in the local universe most of the H I gas with these column densities are in galaxies (e.g. Rosenberg & Schneider 2003; Zwaan et al. 2005). In addition, Prochaska & Wolfe (1998) (and others) argued that the kinematics of DLAs (traced by low-ions) are best explained by models of rapidly rotating disks.

Despite decades of studies, it is not clear whether this cold neutral gas is part of the interstellar medium of large spirals (Wolfe et al. 1986), part of the halos of galaxies (Bahcall & Spitzer 1969) or part of dwarf galaxies (York et al. 1986) akin to the Magellanic Clouds. An alternative scenario for absorption-selected galaxies is that the gas seen in absorption is part of cold gas clumps in the host galaxy halos entrained in outflows produced by supernovae (SNe) (Nulsen et al. 1998; Schaye 2001).

Among the many different approaches that have been used to make further progress on this issue, one is to use the metallicity of the gas. Since the metallicity of the absorbing gas, and of DLAs in particular, is typically 1/30$th$ solar (e.g. Pettini et al. 1999; Prochaska & Wolfe 1999; Péroux et al. 2003; Prochaska et al. 2003), and $z \gg 1$ star-forming galaxies have metallicities closer to solar, i.e. more metal rich by a factor of 10 (e.g. Erb et al. 2006), one is often forced to invoke metallicity gradients in galaxies to 30 kpc and beyond to reconcile the two observations. If indeed absorption and emission metallicities are related to one another solely by a metallicity gradient, one would expect a similar mass-metallicity correlation in DLAs as observed in normal star-forming galaxies (Tremonti et al. 2004; Erb et al. 2006).

While the metallicity of absorbers is usually straightforward to determine (e.g. Prochaska et al. 2000, 2003; Péroux et al. 2003; Péroux et al. 2006; Prochaska et al. 2006; Kulkarni et al. 2007), the mass (dynamical or baryonic) of absorption-selected galaxies cannot be usually measured directly because the host-galaxy is usually elusive. Progress is underway from (i) direct dynamical mass estimates of 14 Mg II-selected galaxies by Bouché et al. (2007) (and Bouché, Murphy & Péroux, in prep.), and (ii) direct comparison between the host-emission metallicity and the absorption metallicity (Bouché et al. 2008, in prep.).

Indirect, and statistical, mass measurements have been made from clustering analysis (Bouché et al. 2006) for strong $z \simeq 0.5$ Mg II absorbers with rest-frame equivalent widths $W_{\rm r}^{\lambda2796} \geqslant 1$ Å, which are intimately related to low-$z$ DLAs (Churchill et al. 2000; Rao & Turnshek 2000; Rao et al. 2006). Bouché et al. (2006)

found that the host halo-mass ($M_h$) and $W_{\rm r}^{\lambda2796}$ are anti-correlated. For virialized clouds, the host mass and the line-of-sight velocity dispersion ought to be correlated, i.e. a $M_h$–$W_{\rm r}^{\lambda2796}$ correlation is expected because $W_{\rm r}^{\lambda2796}$ is a measure of the line-of-sight velocity width ($\Delta v$) as individual Mg II absorptions are saturated (Ellison 2006). Thus, the results of Bouché et al. (2006) imply that Mg II clouds are not virialized in the host halos, and super-novae driven outflows provide a natural mechanism. While the mass-$W_{\rm r}^{\lambda2796}$ results have been confirmed by an independent team (Allen, Hewett & Ryan-Weber 2008, in prep.), ad-hoc models have been proposed to explain the anti-correlation in a cosmological context (e.g. Tinker & Chen 2008).

Since one would expect a mass–metallicity correlation for all galaxies, the mass-$W_{\rm r}^{\lambda2796}$ anti-correlation of Bouché et al. (2006) implies a metallicity-$W_{\rm r}^{\lambda2796}$ anti-correlation, or equivalently a metallicity-velocity width anti-correlation. However, several groups have reported just the opposite: the metallicity in DLAs correlates either with the velocity width $\Delta v$ (Wolfe & Prochaska 1998; Péroux et al. 2003; Ledoux et al. 2006; Prochaska et al. 2008) or with the $W_{\rm r}^{\lambda2796}$ (Meiring et al. 2007; Murphy et al. 2007). It is tempting to assume that $\Delta v$ is a measure of the line-of-sight velocity dispersion, i.e. that it correlates with the mass of the host-galaxy, since when combined with the above results, one would naturally imply a normal mass-metallicity relation.

Thus, there appears to be a conflict between the $M_h$-$W_{\rm r}^{\lambda2796}$ anti-correlation of Bouché et al. (2006) and the $W_{\rm r}^{\lambda2796}$(velocity)–[X/H] correlations reported in the literature. In this paper, we show that the conflict is apparent and reflects the various selections at play (H I vs. Mg II) using Mg II systems from the literature. In section 2, we describe our sample where we collected neutral column density $N_{\rm H\,I}$, Mg II equivalent widths and metallicities [X/H]. Section 3 shows our results.

## 2 DATA

We combined various samples of Mg II absorbers and DLAs from the literature, namely we used the samples of Rao et al. (2006), Ellison (2006), Kulkarni et al. (2005), Curran et al. (2007), Murphy et al. (2007) Ledoux et al. (2006), augmented by the catalog of Ryabinkov et al. (2003). The H I column densities come mostly from the STIS survey of Rao & Turnshek (2000); Rao et al. (2006) for the low-redshift absorbers. The entire catalog contains about 1200 absorbers, of which 377 have both $W_{\rm r}^{\lambda2796}$ and $N_{\rm H\,I}$ measured. We then match the literature samples with published metallicity measurements from Péroux et al. (2003), Péroux et al. (2004), Péroux et al. (2006), Möller et al. (2004). Prochaska et al. (2006), Kulkarni et al. (2005), Ledoux et al. (2006), Ellison (2006), Prochaska et al. (2006), Meiring et al. (2007), Meiring et al. (2008) and Murphy et al. (2007). The final sample is made of 89 absorbers with known $N_{\rm H\,I}$, $W_{\rm r}^{\lambda2796}$, and [X/H].

We show the redshift distribution of the sub-samples in Fig. 1. The solid histogram shows the literature sample of 1200 absorbers. The thick histogram shows the 377 absorbers with $W_{\rm r}^{\lambda2796}$ and $N_{\rm H\,I}$, and the grey histogram shows the 89 absorbers with $W_{\rm r}^{\lambda2796}$, $N_{\rm H\,I}$ and [Zn/H]. In order to have homogeneous metallicity measurements, we impose that all metallicity measurements are from Zn. In order to probe for any redshift evolution, we will split the sample into low-$z$ ($z < 1.6$) and high-$z$ ($z > 1.6$).



**Figure 1.** Redshift distributions of the total sample (solid line), of the H I-Mg II sub-sample (thick line) and of the final sample (shaded histogram) with metallicity measurements restricted to [Zn/H].

## 3 RESULTS

### 3.1 Metallicity gradient in $N_{\rm H\,I}$-$W_{\rm r}^{\lambda2796}$.

Fig. 2(a) shows the distribution of absorbers in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda2796}$ plane as in Rao et al. (2006). Note that strong Mg II absorbers have indeed larger $N_{\rm H\,I}$, and can be regarded as DLA-dominated, however the opposite is not true. A DLA sample covers the entire range of $W_{\rm r}^{\lambda2796}$. The large connected squares show the logarithmic mean $< \log N_{\rm H\,I} >$. The logarithmic mean is a better statistic to quantify the distribution of the points. Rao et al. (2006) elected to use the mean $< N_{\rm H\,I} >$ statistics since they were interested in the mean H I column density in order to constrain $\Omega_{\rm H\,I}$. The solid squares show that $< \log N_{\rm H\,I} >$ increases with equivalent width, reflecting an increasing fraction of DLAs as a function of $W_{\rm r}^{\lambda2796}$.

In Fig. 2(a), as noted many times (e.g. Rao et al. 2006; Chelouche et al. 2007), there are no absorbers to the bottom right of the figure, i.e. with large $W_{\rm r}^{\lambda2796}$ and low H I column densities. The lack of objects in that part of the diagram is not due to selection effects; strong Mg II absorbers are easy to identify there.

The solid squares in Fig. 2(b) show the absorbers whose [X/H] measurement exist in the literature, and color-coded according to [Zn/H] from -2 to 0. This figure clearly shows that the more metal poor (lighter grey points) are located in a different location as the metal rich (darker grey points). In other words, there is a strong metallicity gradient across the $N_{\rm H\,I}$–$W_{\rm r}^{\lambda2796}$ plane, represented schematically in Fig. 3(a). The metallicity gradient is noted by the vector. For low H I column densities ($\log N_{\rm H\,I} < 19.5$), there is a lack of metallicity data due to difficulties in measuring [X/H] due to an increasing ionization correction (Péroux et al. 2006; Prochaska et al. 2006).

### 3.2 Biased metallicities

Fig. 2(b) shows that sub-DLAs are generally more metal rich than DLAs, a result noted already by many (e.g. Khare et al. 2007). However, since the metallicity changes in a subtle way in the $N_{\rm H\,I}$–$W_{\rm r}^{\lambda2796}$ plane, (Fig. 3[a]), one would expect to have different mean metallicity $\langle$[X/H]$\rangle$ for different $N_{\rm H\,I}$-selected samples.



**Figure 2. a)** The distribution of absorbers in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane. Strong Mg II absorbers have indeed larger $N_{\rm H\,I}$, and can be regarded as DLA-dominated, however the opposite is not true. A DLA sample covers the entire range of $W_{\rm r}^{\lambda 2796}$. The large connected squares show that the logarithmic mean $<\log N_{\rm H\,I}>$. **b)** For those with [Zn/H] measurements, the points are color-coded according to increasing metallicity from [Zn/H]=-2 to 0. There is a clear metallicity gradient. Furthermore, the gradient is inclined with respect to the $y$-axis. In both panels, the dashed line shows the classical DLA threshold of 20.3.



**Figure 3. a)**: Schematic representation of the metallicity gradient shown in Fig. 2(b). The 'metal poor' extends over the high $\log N_{\rm H\,I}$ region, while the 'metal rich' region extends over the low $\log N_{\rm H\,I}$ region (assumed to have uniform metallicity). **b)**: Mean metallicity $\langle$[X/H]$\rangle$ for DLAs with $\log N_{\rm H\,I} > 20.3$ (top) and sub-DLAs (bottom). The solid lines show the average $\langle$[X/H]$\rangle$ in bins of $W_{\rm r}^{\lambda 2796}$ as in Fig. 2(a). The increase in the mean metallicity for DLAs is $\sim$ 0.8–1 dex which is the increase reported by Ledoux et al. (2006). On the other hand, sub-DLAs have roughly a similar metallicity. The $P$-values for the Spearman's rank correlation test are shown. This figure shows that the [X/H]-$\Delta v$ and [X/H]-$W_{\rm r}^{\lambda 2796}$ correlations reported by Ledoux et al. (2006) for DLAs and Murphy et al. (2007) are both a 'selection effect': the correlation with metallicity originates from the increased overlap of the metal poor and metal rich systems in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane (Fig. 2b).

This is illustrated in Fig. 3(b), where we plot $\langle$[X/H]$\rangle$ for DLAs with $\log N_{\rm H\,I} > 20.3$ (top) and sub-DLAs with $\log N_{\rm H\,I} < 20.3$ (bottom). The metallicity [X/H] increases as a function of $W_{\rm r}^{\lambda 2796}$ (a proxy for the velocity width $\Delta v$) for DLAs. The $P$-value of the Spearman's correlation test is $2\times 10^{-7}$, i.e. the correlation is significant at $>$ 4-$\sigma$. Moreover, the increase in the mean metallicity [X/H] is $\sim$ 0.8–1 dex, which is the increase reported by Ledoux et al. (2006) for their DLA-sample. On the other hand, for absorbers with $\log N_{\rm H\,I}$ less than 20.3, the mean metallicity appears constant (Fig. 3[b], bottom). The Spearman's correlation test gives a $P$-value much higher (0.10) and shows that the correlation is significant at best at 1.5$\sigma$. Without the one data point at $W_{\rm r}^{\lambda 2796}$=0.5Å, the $P$-value is higher still: 0.43 and the $W_{\rm r}^{\lambda 2796}$ is not correlated with [X/H]. We note that the [X/H]-$W_{\rm r}^{\lambda 2796}$ relation reported by Murphy et al. (2007) is explained by the fact that their sample is dominated by systems with H I column densities mostly above the DLA threshold $N_{\rm H\,I} > 20.3$ (Fig. 2[a]). Thus, the [X/H]-$\Delta v$ and [X/H]-$W_{\rm r}^{\lambda 2796}$ correlations reported by Ledoux et al. (2006) for DLAs and Murphy et al. (2007) are both



**Figure 4.** Based on the metallicity distribution in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane (Fig. 2), we postulate that intervening absorbers in the metal rich region have one physical origin (out-flowing material), whereas absorbers in the metal poor region are 'classical DLAs', in the sense that the sight lines pass through the ISM of galaxies. This simple picture naturally explains the bimodality of DLAs reported by Wolfe et al. (see text).

a 'selection effect': it originates from the increased overlap of the metal poor and metal rich systems in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane.

While the fraction of DLAs increases with $W_{\rm r}^{\lambda 2796}$, this exercise shows that the selection upon $W_{\rm r}^{\lambda 2796}$ is not equivalent to a H I selection, as the two sample will have very different properties: the Mg II-selected sample will be biased towards more metal rich absorbers, while the H I-selected sample will be more metal poor, with a strong metallicity dependence on $W_{\rm r}^{\lambda 2796}$.

Note that there seems to be no redshift bias in our results: Fig. 2[b] looks similar for the $z_{\rm abs} < 1.6$ and $z_{\rm abs} > 1.6$ sub-samples, with perhaps an overall shift in metallicities which has been before for DLAs and sub-DLAs (Prochaska et al. 2003; Kulkarni et al. 2007). Based on these results, we turn towards a physical interpretation.

### 3.3 Interpretation

As we already noted, the $N_{\rm H\,I}$ distribution is bimodal in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane (Fig. 2a). This is particularly true for systems with $W_{\rm r}^{\lambda 2796} < 1.5$ Å. This indicates there might be two classes of intervening absorbers, probing different physical conditions. While observational bias may play a role here, the survey of Rao et al. (2006) is unbiased in regards to H I column density, and the bimodality is already present in their sample.

Given the metallicity gradient shown in the previous section and the absorber distribution in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ plane, we make the following assumption to guide our understanding: absorbers in the red metal-rich shaded region in Fig. 3 originate in one physical environment with a more homogeneous metallicity distribution, while the blue metal-poor absorber in the blue region with low $W_{\rm r}^{\lambda 2796}$and high-$N_{\rm H\,I}$ may be the 'classical' DLAs where the sight-line is probing the ISM of small galaxies. These two hypothesis are illustrated with Fig. 4.

Since our recent results (Bouché et al. 2006, 2007) favor the outflow scenario for Mg II-selected absorbers, we postulate that the homogeneous metallicity distribution originates from the metal-rich material being driven out of sub-L$^*$ galaxies. Indeed, several studies (Bouché et al. 2007; Oppenheimer & Davé 2008) have shown that galaxies with $L < 1/3L^*$ dominate the metal budget in the intergalactic medium, i.e. the metals outside the ISM of galaxies. This material is either directly entrained by ram pressure from the hot outflow, or traces cooling material (where the metallicity is higher and the cooling time is shorter) (Maller & Bullock 2004). Either way it may very well return to the ISM of the galaxy as recent wind models suggest (Oppenheimer & Davé 2008).

This picture outlined in Fig. 4, which may still be somewhat over simplified, naturally explains many other observables of intervening absorbers. For instance, as pointed out in the previous section, a sample of DLAs will be made of a mix of the two types of absorbers. Interestingly, Wolfe et al. (2008) reported evidence for a bimodality in DLAs using [C II] 158$\mu$m cooling rates, $l_c$. The 'low-cool' DLAs have lower velocity widths, lower metallicity and lower dust-to-gas ratios than the 'high-cool' DLAs which have larger velocity widths and higher metallicities. Fig. 3 naturally shows that DLAs with large velocity dispersions (as measured by $W_{\rm r}^{\lambda 2796}$) would be more metal rich, than those with low velocity widths. Furthermore, the UDF results of Wolfe & Chen (2006) imply that in situ star formation can be the dominant heating mechanism for the 'low-cool' population only, consistent with our interpretation shown in Fig. 4. We have shown that the DLA bimodality originates from the biased selection at constant $N_{\rm H\,I}$.

The picture outlined in Fig. 4 predicts an increasing reddening $E(B-V)$ (owing to the [X/H] increase) with equivalent width $W_{\rm r}^{\lambda 2796}$ as reported by York et al. (2006) (see also Ménard et al. 2008) assuming a constant dust-to-gas ratio for Mg II absorbers since globally $W_{\rm r}^{\lambda 2796}$ and $N_{\rm H\,I}$ are correlated. This assumption may be the case under the common physical nature (out-flowing material) of strong Mg II samples. After this work was being completed, Ménard & Chelouche (2008) showed that Mg II absorbers on the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ sequence shown in Fig. 2(a) have indeed a constant dust-to-gas ratio Interestingly, they concluded that the dust-to-gas ratio was not consistent with that of dwarf galaxies (SMC), therefore rejecting the alternative hypothesis often invoked for absorbers, namely that the sight-lines go through dwarfs near more normal galaxies. We note that the dust-to-gas ratio for the data points in the upper left of Fig. 2(a) (i.e. not on $N_{\rm H\,I}$-$W_{\rm r}^{\lambda 2796}$ sequence) must be smaller (Ménard & Chelouche 2008), as concluded by Wolfe et al. (2008) for DLAs with low velocity widths.

### 3.4 Caveats and possible systematics

Both Prochaska et al. (2008) and Ledoux et al. (2006) used a sample of DLAs at higher redshifts (with $1.7 < z_{\rm abs} < 4.3$) than the Rao et al. sample of low-$z$ absorbers dominating our study. However, Fig. 2(b) does not change qualitatively for absorbers with $z_{\rm abs} > 1.6$ in our sample. The overall metallicity is lower, reflecting the redshift evolution of metallicity in DLAs and sub-DLAs (Prochaska et al. 2003; Péroux et al. 2007; Kulkarni et al. 2007).

Another difference between our analysis and Ledoux et al. (2006) and Prochaska et al. (2008), is that we used $W_{\rm r}^{\lambda 2796}$ as a proxy for velocity width $\Delta v$ (Ellison 2006) whereas they used the measured velocity width from high-resolution spectra of the low-ions Si II and Zn II. However, for the dozen of absorbers with both $W_{\rm r}^{\lambda 2796}$ and $\Delta v(\text{Si II or Zn II})$, we find that the $\Delta v(W_{\rm r}^{\lambda 2796})$ correlates with $\Delta v(\text{Si II or Zn II})$ at 99% confidence level. Thus, the selection effect shown in Fig. 3 exists against $W_{\rm r}^{\lambda 2796}$ (as in Murphy et al. 2007) and against $\Delta v(\text{Si II or Zn II})$ (as in Ledoux et al. 2006; Prochaska et al. 2008).

One might invoke dust obscuration to account for the results shown in Fig. 3. However, such a dust-bias would have to selectively remove metal rich DLAs with low $W_{\rm r}^{\lambda2796}$. The results of York et al. (2006) and Ménard et al. (2008) showed that dust obscuration in Mg II-selected samples with the lowest $W_{\rm r}^{\lambda2796}$ is low, i.e. $E(B-V) < 0.02$ below $W_{\rm r}^{\lambda2796} < 2$ Å, which implies a very low fraction of missed absorbers below $W_{\rm r}^{\lambda2796} < 2$ Å irrespective of $N_{\rm H\,I}$. Similarly, such a dust-bias would have to selectively remove metal poor DLAs with high $W_{\rm r}^{\lambda2796}$. This is unlikely since dusty and metal poor ([X/H]<-1.0) DLAs would have to have unphysically high dust-to-metal ratios ($\mathcal{D} \geqslant 10\mathcal{D}_{\rm MW}$), which is rather difficult to produce (Inoue 2003). On the other hand, metal-rich DLAs with high $W_{\rm r}^{\lambda2796}$ would have larger $E(B-V)$ (Ménard & Chelouche 2008) and therefore would be easier to obscure, but are in fact present in our sample. Finally, we looked at the distribution of the QSO magnitudes in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda2796}$ plane, and found no evidence for a selective dust-bias.

## 4 CONCLUSIONS

Our results show the presence of a metallicity gradient in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda2796}$ plane of intervening absorbers (Fig. 2). In other words, the metallicity of absorbers is a bimodal function of $N_{\rm H\,I}$ and $W_{\rm r}^{\lambda2796}$. As a direct consequence, a population of DLAs, selected with $\log N_{\rm H\,I} > 20.3$, will be heterogeneous. At low $W_{\rm r}^{\lambda2796}$, the H I-selected sample is metal poor, whereas at high $W_{\rm r}^{\lambda2796}$, it is found to be more metal rich. Therefore, the correlation between the metallicity [X/H] and the line-of-sight velocity width $\Delta v$ reported by Ledoux et al. (2006) and by Murphy et al. (2007) arises from the H I selection and can not be interpreted as a signature of the mass-metallicity relation akin to normal field galaxies.

We argue that the bivariate distribution [[X/H]($N_{\rm H\,I}$,$W_{\rm r}^{\lambda2796}$)] can be explained by two different physical origins of absorbers, which are the ISM of small galaxies and out-flowing material, with distinct physical properties (such as metallicities and dust-to-gas ratios). This is supported by the distribution of the absorbers in the $N_{\rm H\,I}$-$W_{\rm r}^{\lambda2796}$ plane. If there are two distinct populations of absorbers, as shown in Fig. 4, this naturally explains the two classes of DLAs ('low-cool' and 'high-cool'), reported by Wolfe et al. (2008) using [C II] 158$\mu$m cooling rates. DLAs with large velocity dispersions (as measured by $W_{\rm r}^{\lambda2796}$) are more metal rich than those with low velocity widths and will have different dust-to-gas ratio for a given dust-to-metal ratio.

Therefore, the correlation between metallicity [X/H] and $W_{\rm r}^{\lambda2796}$(or $\Delta v$) for DLAs (which we showed to be apparent), the two classes of DLAs of Wolfe et al. (2008), the dust-to-gas results of Ménard & Chelouche (2008), and the results of Bouché et al. (2006) indicating that Mg II-selected absorbers are tracing out-flowing material can all be put into one coherent context.

## ACKNOWLEDGMENTS

We thank S. Ellison for providing $W_{\rm r}^{\lambda2796}$ for her sample. We acknowledge inlighting scientific discussions with M.T. Muprhy, C. Péroux. We gratefully acknowledge M.T. Murphy and S. Genel for their thorough reading of the manuscript.

## REFERENCES

Bahcall J. N., Spitzer L. J., 1969, ApJ, 156, L63+
Bouché N., Lehnert M. D., Aguirre A., Péroux C., Bergeron J., 2007, MNRAS, 378, 525
Bouché N., Lehnert M. D., Péroux C., 2006, MNRAS, 367, L16
Bouché N., Murphy M. T., Peroux C., Csabai I., Wild V., 2006, MNRAS, 371, 495
Bouché N., Murphy M. T., Péroux C., Davies R., Eisenhauer F., Förster Schreiber N. M., Tacconi L., 2007, ApJ, 669, L5
Chelouche D., Ménard B., Bowen D. V., Gnat O., 2007, ApJ, (astro-ph/0706.4336), 706
Churchill C. W., Mellon R. R., Charlton J. C., Jannuzi B. T., Kirhakos S., Steidel C. C., Schneider D. P., 2000, ApJ, 543, 577
Curran S. J., Tzanavaris P., Pihlström Y. M., Webb J. K., 2007, MNRAS, 382, 1331
Ellison S. L., 2006, MNRAS, 368, 335
Erb D. K., Shapley A. E., Pettini M., Steidel C. C., Reddy N. A., Adelberger K. L., 2006, ApJ, 644, 813
Inoue A. K., 2003, Publ. Astron. Soc. Japan, Vol.55, No.5, pp. 901-909, 55, 901
Khare P., Kulkarni V. P., Péroux C., York D. G., Lauroesch J. T., Meiring J. D., 2007, A&A, 464, 487
Kulkarni V. P., Fall S. M., Lauroesch J. T., York D. G., Welty D. E., Khare P., Truran J. W., 2005, ApJ, 618, 68
Kulkarni V. P., Khare P., Péroux C., York D. G., Lauroesch J. T., Meiring J. D., 2007, ApJ, 661, 88
Ledoux C., Petitjean P., Fynbo J. P. U., Møller P., Srianand R., 2006, A&A, 457, 71
Meiring J. D., Kulkarni V. P., Lauroesch J. T., Péroux C., Khare P., York D. G., Crotts A. P. S., 2008, MNRAS, 384, 1015
Maller, A. H. and Bullock, J. S. 2004, MNRAS, 355, 694
Meiring J. D., Lauroesch J. T., Kulkarni V. P., Péroux C., Khare P., York D. G., Crotts A. P. S., 2007, MNRAS, pp 112–+
Ménard B., Chelouche D., 2008, MNRAS, (astro-ph/0803.0745)
Ménard B., Nestor D., Turnshek D., Quider A., Richards G., Chelouche D., Rao S., 2008, MNRAS, pp 219–+
Möller P., Fynbo J. P. U., Fall S. M., 2004, A&A, 422, L33
Murphy M. T., Curran S. J., Webb J. K., Ménager H., Zych B. J., 2007, MNRAS, pp 74–+
Nulsen P. E. J., Barcons X., Fabian A. C., 1998, MNRAS, 301, 168
Oppenheimer B. D., Davé R., 2008, ArXiv e-prints, 712
Péroux C., Deharveng J.-M., Le Brun V., Cristiani S., 2004, MNRAS, 352, 1291
Péroux C., Dessauges-Zavadsky M., D'Odorico S., Kim T., McMahon R. G., 2003, MNRAS, 345, 480
Péroux C., McMahon R. G., Storrie-Lombardi L. J., Irwin M. J., 2003, MNRAS, 346, 1103
Péroux C., Dessauges-Zavadsky M., D'Odorico S., Kim T.-S., McMahon R. G., 2007, MNRAS, 382, 177
Péroux C., Kulkarni V. P., Meiring J., Ferlet R., Khare P., Lauroesch J. T., Vladilo G., York D. G., 2006, A&A, 450, 53
Pettini M., Ellison S. L., Steidel C. C., Bowen D. V., 1999, ApJ, 510, 576
Prochaska J. X., Chen H.-W., Wolfe A. M., Dessauges-Zavadsky M., Bloom J. S., 2008, ApJ, 672, 59
Prochaska J. X., Gawiser E., Wolfe A. M., Castro S., Djorgovski S. G., 2003, ApJ, 595, L9
Prochaska J. X., Gawiser E., Wolfe A. M., Cooke J., Gelino D., 2003, ApJS, 147, 227
Prochaska J. X., O'Meara J. M., Herbert-Fort S., Burles S., Prochter G. E., Bernstein R. A., 2006, ApJ, 648, L97
Prochaska J. X., Wolfe A. M., 1998, ApJ, 507, 113
Prochaska J. X., Wolfe A. M., 1999, ApJS, 121, 369
Prochaska J. X., Wolfe A. M., Gawiser E. J., 2000, BAAS, 32, 3.09
Rao S. M., Turnshek D. A., 2000, ApJS, 130, 1
Rao S. M., Turnshek D. A., Nestor D. B., 2006, ApJ, 636, 610
Rosenberg J. L., Schneider S. E., 2003, ApJ, 585, 256
Ryabinkov A. I., Kaminker A. D., Varshalovich D. A., 2003, A&A, 412, 707
Schaye J., 2001, ApJ, 559, L1
Tinker J. L., Chen H.-W., 2008, ApJ, astro-ph/0709.1470, 709

Tremonti C. A., Heckman T. M., Kauffmann G., Brinchmann J., Charlot S., White S. D. M., Seibert M., Peng E. W., Schlegel D. J., Uomoto A., Fukugita M., Brinkmann J., 2004, ApJ, 613, 898
Wolfe A. M., Chen H.-W., 2006, ApJ, 652, 981
Wolfe A. M., Prochaska J. X., 1998, ApJ, 494, L15+
Wolfe A. M., Prochaska J. X., Jorgenson R. A., Rafelski M., 2008, ArXiv e-prints, 802
Wolfe A. M., Turnshek D. A., Smith H. E., Cohen R. D., 1986, ApJS, 61, 249
York D. G., Burks G. S., Gibney T. B., 1986, AJ, 91, 354
York D. G., et al., 2006, MNRAS, 367, 945
Zwaan M. A., van der Hulst J. M., Briggs F. H., Verheijen M. A. W., Ryan-Weber E. V., 2005, MNRAS, 364, 1467

This paper has been typeset from a TEX/ LATEX file prepared by the author.